

Control of gluconeogenic genes during intense/prolonged exercise: hormone-independent effect of muscle-derived IL-6 on hepatic tissue and PEPCK mRNA.

Sébastien Banzet, Nathalie Koulmann, Nadine Simler, Hervé Sanchez, Rachel Chapot, Bernard Serrurier, André Peinnequin, Xavier Bigard

► To cite this version:

Sébastien Banzet, Nathalie Koulmann, Nadine Simler, Hervé Sanchez, Rachel Chapot, et al.. Control of gluconeogenic genes during intense/prolonged exercise: hormone-independent effect of muscle-derived IL-6 on hepatic tissue and PEPCK mRNA.. *Journal of Applied Physiology*, American Physiological Society, 2009, 107 (6), pp.1830-9. <10.1152/jappphysiol.00739.2009>. <ssa-00439585>

HAL Id: ssa-00439585

<https://hal-ssa.archives-ouvertes.fr/ssa-00439585>

Submitted on 7 Dec 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1

2 Control of gluconeogenic genes during intense/prolonged
3 exercise: hormone independent effect of muscle-derived IL-6
4 on hepatic tissue and PEPCK mRNA

5

6 Banzet Sébastien, Koulmann Nathalie, Simler Nadine, Sanchez Hervé, Chapot Rachel,
7 Serrurier Bernard, Peinnequin André and Bigard Xavier.

8 Department of Human Factors,

9 Military health service research center (Centre de Recherches du Service de Santé des

10 Armées),

11 38702 La Tronche, France

12

13 Running head : IL-6 and exercise gluconeogenesis

14

15 **Address for correspondence:**

16 Sébastien Banzet

17 CRSSA

18 Department of Human Factors

19 BP 87

20 38702 La Tronche Cedex

21 France

22

23 Phone : +334 76 63 97 67

24 Fax : +334 76 63 69 45

25 E-mail: sbanzet@crssa.net

26

Abstract

27
28 Prolonged intense exercise is challenging for the liver to maintain plasma glucose levels.
29 Hormonal changes cannot fully account for exercise-induced hepatic glucose production
30 (HGP). Contracting skeletal muscles release interleukin-6 (IL-6), a cytokine able to increase
31 endogenous glucose production during exercise. However, whether this is due to a direct
32 effect of IL-6 on liver remains unknown. Here, we studied hepatic glycogen, gluconeogenic
33 genes and IL-6 signaling in response to one bout of exhaustive running exercise in rats. To
34 determine whether IL-6 can modulate gluconeogenic genes mRNA independently of exercise,
35 we injected resting rats with recombinant IL-6. Exhaustive exercise resulted in a profound
36 decrease in liver glycogen and an increase in gluconeogenic genes mRNA levels,
37 phosphoenolpyruvate-carboxykinase (PEPCK), glucose-6-phosphatase (G6P) and peroxisome
38 proliferator-activated receptor γ coactivator-1 α (PGC-1 α), suggesting a key role for
39 gluconeogenesis in hepatic glucose production. This was associated to an active IL-6
40 signaling in liver tissue, as shown by signal transducer and activator of transcription (STAT-
41 3) and CAAT/enhancer binding protein- β (C/EBP β) phosphorylation and IL-6 responsive
42 genes mRNA levels at the end of exercise. Recombinant IL-6 injection resulted in an increase
43 in IL-6 responsive genes mRNA levels in the liver. We found a dose-dependent increase in
44 PEPCK gene mRNA, strongly correlated with IL-6-induced genes mRNA levels. No changes
45 in G6P and PGC-1 α mRNA levels were found. Taken together, our results suggest that during
46 very demanding exercise, muscle-derived IL-6 could help increasing HGP by directly up
47 regulating PEPCK mRNA abundance.

48
49

50 During intense and prolonged exercise, working skeletal muscles are very dependent
51 on carbohydrates to maintain ATP synthesis and contraction (37). In the absence of oral
52 glucose intake, plasma glucose levels are maintained at relatively constant values through
53 changes in hepatic glucose production (HGP). During moderate-intensity exercise, HGP is
54 primarily controlled by portal venous insulin-to-glucagon ratio (44, 45), whereas during
55 intense and sustained exercise, several studies have shown that hormonal changes cannot fully
56 account for the increase in HGP, suggesting other unidentified factors are involved (9, 20). A
57 new and original aspect of HGP control during intense exercise as been highlighted recently
58 by a vast series of studies supporting the involvement of muscle-derived factor (for review see
59 (32)).

60 Contracting skeletal muscles produce and release IL-6 in the plasma during prolonged
61 sustained exercise (30, 39), and a strong relationship with low muscle glycogen content as
62 been shown (25, 38). It has been proposed that this cytokine may be a signal of low muscle
63 glycogen stores and, once released in the plasma, could favor hepatic glucose release (13).
64 This hypothesis has been tested in an experiment were subjects exercised with an infusion of
65 saline or recombinant IL-6. The rate of glucose appearance was significantly higher in the IL-
66 6 group with no changes in gluco-regulatory hormones, suggesting IL-6 is involved in HGP
67 increase during intense prolonged exercise (11). However, the mechanisms involved are
68 unknown and the question was asked whether this effect was due to a direct effect of IL-6 on
69 hepatic tissue or was secondary to IL-6-induced increase in whole-body glucose disposal (11).

70

71 Exercise-induced increase in HGP results from coordinated changes in glycogenolysis
72 and gluconeogenesis (GNG) under complex hormonal and nervous control (8, 26). Liver
73 glycogenolysis starts at the onset of exercise and is the main source of glucose during the first
74 30 min of exercise (26). Thereafter, hepatic gluconeogenic flux increases and GNG becomes

75 a key contributor to HGP (44). GNG is controlled at the transcriptional level by unidirectional
76 key enzymes, phosphoenolpyruvate-carboxykinase (PEPCK), fructose 1,6 bisphosphatase and
77 glucose-6 phosphatase (G6Pase) the activities of which increase during exercise (10).
78 PEPCK, the most studied gluconeogenic enzyme, plays an important role during exercise
79 since its pharmacologic inhibition reduces endurance time in rats (22, 42). PEPCK activity is
80 hormonally controlled at the transcriptional level (16) and exercise induces an important
81 increase in hepatic PEPCK mRNA levels and enzyme activity (14, 44). Among the well-
82 described regions of PEPCK gene promoter, the glucocorticoid response unit (GRU) and two
83 cAMP regulatory element (CRE) are involved in exercise-related gene expression, since their
84 mutation substantially decreased hepatic PEPCK mRNA levels (14, 29). However, the
85 molecular mechanisms controlling PEPCK gene expression in response to exercise is still not
86 fully understood. Recently, peroxisome proliferator-activated receptor γ coactivator-1 α (PGC-
87 1 α) has been shown to play a pivotal role in hepatic GNG regulation and PEPCK gene
88 transcription in response to food deprivation (33, 46). An increase in PGC-1 α mRNA levels
89 has been reported recently in exercised mice (19), however its role in hepatic GNG is
90 unknown.

91 IL-6 binds to a specific receptor, IL-6 receptor α (IL-6R α) that cooperates with a non-
92 specific subunit, glycoprotein 130 (gp130). The signal transduction occurs through either the
93 janus kinase (JAK)/signal transducer and activator of transcription (STAT) pathway, or a
94 Ras/extracellular signal-regulated protein kinase (ERK)/CAAT enhancer binding protein
95 (C/EBP) β pathway (for review see (23)). C/EBP family members can trans activate PEPCK
96 promoter via the CRE binding sites (16, 31). C/EBP β , also known as nuclear factor of IL-6
97 (NF-IL6), has been proposed to play a role in PEPCK gene transcription during exercise (29).
98 Because this transcription factor is regulated by IL-6 at the transcriptional level and is a
99 downstream signaling molecule of activated IL-6 receptor (reviews in (1, 35)), muscle-

100 derived IL-6 could participate in hepatic C/EBP β activation during exercise. Whether IL-6
101 can act directly on hepatic tissue to favor PEPCK gene transcription during exercise has to be
102 explored.

103 The present study focused on prolonged and intense exercise to further understand
104 glucose homeostasis in very demanding physiologic situations. Because plasma IL-6
105 concentration is a function of exercise intensity and duration (32) and because muscle-IL-6
106 release is concomitant of GNG increase, we hypothesized that this cytokine could act directly
107 on hepatic tissue to help increasing HGP through GNG. Our aims were 1) to determine if
108 there is an active IL-6 signaling in the liver of animals exposed to one bout of prolonged
109 intense exercise; 2) to determine if IL-6, can directly modulate gluconeogenic genes mRNA
110 abundance, independently of exercise and glucoregulatory hormones. Therefore, we studied
111 liver IL-6 signaling and gluconeogenic genes mRNAs at the end of one bout of exhaustive
112 exercise in rats. To evaluate the ability of IL-6 to directly activate gluconeogenic genes
113 independently of exercise, we studied gluconeogenic genes modulation in response to
114 recombinant IL-6 injection in resting animals.

115

116 **Research Design and Methods**

117

118 **Ethics**

119 Experiments were carried out in accordance with the Helsinki Accords for Human Treatment
120 of Animals during Experimentation, and received prior approval from the animal ethics
121 committee for animal research of the Centre de Recherche du Service Santé des Armées (La
122 Tronche, France).

123

124 **Animals**

125 Female Wistar rats (175-200 g) were purchased from Charles River (L'Arbresle, France).
126 They were housed at $22 \pm 2^{\circ}\text{C}$, on a 12–12 h light–dark period, and provided with food and
127 water *ad libitum*. For the exercise experiment they were randomly assigned to four
128 experimental groups: resting rats (Rest, $n=8$) and 3 active groups, performing one bout of
129 exercise and studied at the end of exercise (Ex, $n=8$), 2 hours (Ex+2h, $n=8$) or 6 hours
130 (Ex+6h, $n=8$) later.

131

132 **Exercise protocol**

133 All animals were accustomed to running on a rodent treadmill for 15-20 min per day for 5
134 days ($10\text{--}20 \text{ m}\cdot\text{min}^{-1}$; 0° grade). After 4 days at rest, active animals ran on the treadmill (22
135 $\text{m}\cdot\text{min}^{-1}$, 5° grade) till exhaustion.

136

137 **IL-6 injection**

138 Animals were fasted overnight and injected intraperitoneally with 3 or $10 \mu\text{g}\cdot\text{kg body weight}^{-1}$
139 human recombinant IL-6 (Biosource, Camarillo, CA, USA) and compared to control rats
140 injected with endotoxin free PBS (Cont). Animals were sacrificed 1 hour after injection.

141

142 **Tissue processing**

143 Resting animals were sacrificed at different time corresponding to the other groups. After
144 exercise, rats from Ex+2h and Ex+6h groups had free access to food and water. Animals were
145 anaesthetized with pentobarbital (70 mg.kg body weight⁻¹ *ip*). Liver was sampled and a part
146 was frozen in liquid nitrogen and stored at -80°C for protein and glycogen quantification,
147 another part was placed in 1ml RNAlater (Ambion, Austin, TX, USA), kept at 4°C for 24 h
148 and then frozen at -20°C for mRNA study. Whole blood was sampled, centrifuged and
149 plasma was frozen at -80°C. Animals were killed by removal of the heart.

150

151 **RNA extraction**

152 Hepatic tissue mRNA were isolated using the MagNA Pure LC instrument (Roche Applied
153 Science, Mannheim, Germany). 150 to 200 mg of liver were disrupted in a volume
154 corresponding to 6,66 µl MagNA Pure LC mRNA isolation kit II lysis buffer (Roche Applied
155 Science, Mannheim, Germany) per mg of tissue, with a Polytron mixer (Kinematica,
156 Switzerland). Lysate was centrifuged and 50µl was transferred in a fresh tube and completed
157 to a 300-µl final volume with lysis buffer. Extraction was then performed following the
158 manufacturer's protocol.

159

160 **Reverse transcription**

161 Reverse transcription was carried out using the Reverse Transcriptase Core Kit (Eurogentec,
162 Seraing, Belgium). The reaction was performed following the manufacturer's instructions.
163 The reaction mix contained 1 µl buffer (10X), 2 µl dNTP (2.5 mM each dNTP), 0.5 µl oligo-
164 dT primer (50 µM), 0.2 µl RNase inhibitor (20 units.µl⁻¹), 2 µl MgCl₂ (25 mM), 0.25 µl
165 Euroscript reverse transcriptase (50 U.µl⁻¹), 1.05 µl RNase free water and 3 µl template
166 mRNA.

167

168 **Real-time PCR**

169 The PCR reactions were carried out in a 20 μ l final volume with the LC Fast Start DNA
170 Master SYBR Green kit (Roche Applied Science, Mannheim, Germany) using 2 μ l of cDNA
171 solution (1/20 dilution). Quantitative PCRs were performed using (Roche Applied Science,
172 Mannheim, Germany) with 50 cycles of 95 °C for 20 s, annealing temperature (table 1) for 5 s
173 and 72 °C for 8 s. Quantification was performed with the comparative threshold cycle method
174 (28) using RelQuant software (Roche Diagnostics), with a pool of cDNA samples as
175 calibrator. Five housekeeping genes were necessary to validate the normalization using
176 genorm software (43). These were Hypoxanthine-guanine phosphoribosyl transferase
177 (HPRT), cyclophilin A (CycA), acidic ribosomal binding protein (ARBP), actin β and
178 glyceraldehydes-3-phosphate dehydrogenase (GAPDH). Primers characteristics are provided
179 as supplemental data. The final quantification was performed using the geometric averaging
180 of the five quantifications. In each experiment, all samples were treated at the same time for
181 mRNA extraction, reverse transcription and PCR reaction.

182

183 **Western blotting**

184 Liver sections were homogenized in ice-cold lysis buffer (20 μ l.mg⁻¹) containing 20 mM
185 HEPES, pH 7.4, 2 mM EGTA, 50 mM β -glycerophosphate, 1 mM dithiothreitol, 1 mM
186 Na₃VO₄, 1% Triton X-100, 10% glycerol, 1 μ l.ml⁻¹ Protease inhibitor cocktail set III
187 (Calbiochem, Fontenay-sous-bois, France). Samples were rotated for 1 h at 4°C and
188 centrifuged (15,000 g, 15 min, 4°C). Protein content was determined using the BCA method
189 (Roche/Hitachi 912 instrument, Roche Diagnostics, Mannheim, Germany).

190 Each sample (50 μ g of total proteins) was separated by SDS-PAGE (10% resolving gel).
191 Proteins were transferred electrophoretically on nitrocellulose membranes (Hybond C-extra
192 RPN 2020E, Amersham, Orsay, France). Membranes were blocked with non-fat milk in Tris-
193 buffered-Saline solution added with 0.1% Tween 20 (TBST). Blots were incubated overnight
194 at 4°C with anti-phospho-STAT-3 (Ser727) or anti-phospho-C/EBP β (Ser105) antibody (Cell

195 Signaling Technology, Beverly, MA, USA), rinsed with TBST and incubated with horse
196 radish peroxidase-conjugated donkey anti-rabbit antibody (Santa Cruz Biotechnology, Santa
197 Cruz, CA, USA) for 1h30. Blots were subjected to Enhanced chemiluminescence reagent kit
198 (Amersham, Orsay, France) and exposed to Hyperfilm ECL RPN 3103K (Amersham, Orsay,
199 France). Membranes were then stripped in a solution containing 60 mM Tris-HCL, pH 7.4,
200 5% SDS and 0.07% β -mercapto ethanol, for 30 min at 50°C, washed with TBST, and
201 incubated overnight with anti STAT-3 antibody or C/EBP β (Cell Signaling Technology,
202 Beverly, MA, USA) and treated as described above. The bands obtained were quantified
203 using a densitometer (GS-700, Bio-Rad, Marnes-la-Coquette, France).

204

205 **Liver and muscle glycogen content**

206 Liver or plantaris muscle sections (30 mg or 20 mg respectively) were disrupted in NaOH (2
207 M). Samples were submitted to amyloglucosidase (Sigma-Aldrich, Steinheim, Germany) in
208 acetate buffer (0.3 M) for 2 hours at 37°C to digest glycogen. Released glucose was
209 quantified by spectrophometric measurement (340 nm) of NADH production in the presence
210 of hexokinase and glucose-6-phosphate dehydrogenase (Sigma-Aldrich, Steinheim,
211 Germany), as previously described (4).

212

213 **Blood glucose**

214 Blood glucose was measured according to the glucose oxidase methods with Glucose GOP-
215 PAP kit on Roche/Hitachi 912 instrument (Roche Diagnostics, Mannheim, Germany).

216

217 **IL-6 protein**

218 Plasma IL-6 levels were measured using the Quantikine colorimetric sandwich ELISA kit for
219 rat IL-6 (R&D systems, Lille, France) in exercise experimentation and sandwich ELISA kit
220 for human IL-6 (R&D systems, Lille, France) in recombinant IL-6 injected rats.

221

222 **Hormones**

223 Plasma hormones were measured using BioPlex Instrument (Biorad, France) with rat endo kit
224 (Millipore, France) for glucagon and insulin, rat stress kit for corticosterone.

225

226 **Statistical analysis**

227 All data are presented as means \pm S.E.M. A one factor ANOVA was used to evaluate the
228 global effects of exercise and recovery, as well as recombinant IL-6 injection. When
229 appropriate, the Newman-Keuls *post hoc* test was used for inter-group comparisons. Animals
230 weight and running time values were compared using an unpaired Student *t* test. The
231 significance level for all comparisons was set at $P < 0.05$.

232

233

234 **Results**

235 **Liver response to exercise**

236 *Endurance time:* Mean endurance time was 123±10 min with no differences between groups
237 (data not shown).

238
239 *Carbohydrate metabolism:* Plasma glucose was lower in Ex group than Rest and Ex+2h (8.25
240 ± 0.33; 6.71 ± 0.51 and 8.26 ± 0.34 mmol.l⁻¹ respectively, data not shown) ($P<0.05$). Plantaris
241 muscle glycogen content decreased immediately at the end of exercise (Rest: 24.11 ± 0.74;
242 Ex: 3.37 ± 0.6 μmol glycosyl units.g⁻¹, data not shown) ($P<0.01$). There was a profound
243 decrease in hepatic glycogen in response to exercise (main effect $P<0.001$), with only 5% of
244 the resting values at the end of exercise ($P<0.001$). Liver glycogen increased during recovery,
245 but was still 35% lower in Ex+6h than in Rest group ($P<0.001$) (fig 1A).

246
247 *Gluconeogenic genes:* Prolonged running exercise induced an increase in both PEPCK and
248 G6Pase mRNA levels (main effects $P<0.001$) with a maximum at the end of exercise
249 ($P<0.001$). Mean mRNA values decreased to values measured in Rest group as soon as 2h
250 after exercise (fig 1B and 1C). Changes in PGC-1α mRNA levels exhibited a similar
251 evolution ($P<0.0001$) (fig 1D).

252
253 *Glucoregulatory hormones:* Exercise resulted in a prolonged decrease in plasma
254 insulin/glucagon ratio (main effect $P<0.01$), values being lower than control in Ex and Ex+2h
255 groups ($P<0.05$) (fig 2A). Plasma corticosterone concentrations increased in response to
256 running (main effect $P<0.05$), peaking at the end of exercise (fig 2B). Concentrations
257 increased 6h after exercise ($P<0.05$) (fig 2B).

258
259 *Plasma IL-6:* Exercise induced an increase in plasma IL-6 protein (main effect $P<0.05$)
260 peaking at the end of exercise ($P<0.01$). Plasma IL-6 concentrations decreased to values
261 measured in resting animals as soon as 2h after exercise (fig 3A).

262
263 *IL-6 signaling in liver tissue:* Liver STAT-3 and C/EBPβ phosphorylation ratio significantly
264 increased in response to exercise (global effects $P<0.05$) (fig 3B and C), with a maximum at
265 the end of exercise ($P<0.05$ and $P<0.01$ respectively).

266 Hepatic mRNA levels for SOCS-3 and C/EBP β , two IL-6 responsive genes, substantially
267 increased (main effects $P<0.001$) peaking at the end of exercise ($P<0.001$) (fig 4A and B).
268 SOCS-3 mRNA levels were still slightly elevated at Ex+2h compared to resting values
269 ($P<0.05$). Interestingly, we found a 3.5 fold increase in IL-6R mRNA at the end of exercise
270 ($P<0.05$) (fig 4C).

271

272 *Correlation between PEPCK and IL-6-induced genes mRNAs in Rest an Exercise groups:* We
273 found significant correlations between PEPCK mRNA levels and two IL-6-induced genes
274 mRNA levels, SOCS-3 ($R^2= 0.5546$; $P<0.05$) (fig 7A) and IL-6R ($R^2 = 0.7707$; $P<0.01$) (fig
275 7B).

276

277 **Liver response to recombinant IL-6**

278 *IL-6 responsive genes in liver tissue:* Recombinant IL-6 injection resulted in a dose dependent
279 peak of recombinant IL-6 plasma levels (fig 5A) ($P<0.01$) and induced an increase in both
280 SOCS-3 and IL-6R mRNA levels ($P<0.01$ and $P<0.05$ respectively) (fig 5B and C).

281

282 *Gluconeogenic genes:* Recombinant IL-6 injection induced a dose dependent increase in
283 hepatic PEPCK mRNA levels (main effect $P<0.01$) (fig 6A). No effects of IL-6 injection was
284 found on G6Pase and PGC-1 α mRNA levels (fig 6B and C).

285

286 *Plasma glucose and gluco-regulatory hormones:* Recombinant IL-6 had no effect on plasma
287 glucose (data not shown). There were no changes in plasma insulin/glucagon ratio ($15.07 \pm$
288 3.42 , 13.03 ± 2.94 and 14.03 ± 1.69 for Cont, $3 \mu\text{g.kg}^{-1}$ and $10 \mu\text{g.kg}^{-1}$ groups, respectively)
289 and corticosterone was unchanged ($252.27 \pm 38.24 \text{ ng.ml}^{-1}$, $254.81 \pm 35.12 \text{ ng.ml}^{-1}$ and
290 $247.48 \pm 49.16 \text{ ng.ml}^{-1}$ for Cont, $3 \mu\text{g.kg}^{-1}$ and $10 \mu\text{g.kg}^{-1}$ groups, respectively).

291

292 *Correlation between PEPCK and IL-6-induced genes mRNAs:* We found significant
293 correlations between PEPCK mRNA levels and two IL-6-induced genes mRNA levels,
294 SOCS-3 ($R^2= 0.5764$; $P<0.01$) (fig 7C) and IL-6R ($R^2 = 0.6094$; $P<0.01$) (fig 7D).

295

296

297 **Discussion**

298 During prolonged intense exercise, a strong relationship between muscle IL-6
299 production and carbohydrate availability has been described (32) and IL-6 is able to increase
300 endogenous glucose production in exercising humans (11). However, the mechanisms
301 involved in IL-6 action are unknown. In this work, we show that: 1) prolonged intense
302 exercise is associated to an active IL-6 signaling in the liver, involving both STAT-3 and
303 C/EBP β pathways; 2) Recombinant IL-6 can acutely modulate hepatic PEPCK mRNA levels
304 independent from exercise and gluco-regulatory hormones.

305 We previously reported that exhaustive running exercise protocol in rats induces an
306 increase in muscle IL-6 transcription and in plasma IL-6 concentration (3). This was
307 confirmed in the present study where plasma IL-6 peaked at the end of exercise and decreased
308 as soon as 2 hours recovery, as described in human (30). However, whether such a transient
309 increase in circulating IL-6 can induce an active IL-6 signaling in the liver has never been
310 assessed. When IL-6 binds to its receptor, signal transduction can involve STAT-3 and/or
311 C/EBP β transcription factors to promote gene transcription (1). In the present study we show
312 that in addition to increased hepatic C/EBP β phosphorylation, STAT-3 phosphorylation
313 substantially raised at the end of exercise. Therefore, plasma IL-6 concentration and IL-6-
314 activated transcription factors phosphorylation followed the same time course. When
315 phosphorylated, both transcription factors are translocated in the nucleus (1), and to determine
316 if transcription factors activation was effective in altering transcription in our experiment, we
317 measured hepatic mRNA levels for IL-6-induced genes. SOCS-3, a cytokine signaling
318 suppressor molecule, is highly induced at the transcriptional level by IL-6 (15). Here, we
319 found an important increase in liver SOCS-3 mRNA in response to exercise, still significant
320 after 2 hours recovery. Although commonly used to assess IL-6 activity, SOCS-3 is not fully

321 specific of IL-6 and we studied IL-6R and C/EBP β , two IL-6-inducible genes in the liver in
322 vivo (2, 36). We found that mRNA levels for both genes were transiently activated at the end
323 of exercise. Taken together, concomitant elevation in plasma IL-6, the increase in its
324 transcription factors phosphorylation and IL-6-responsive genes activation in the liver
325 strongly suggests that prolonged intense exercise can acutely induce an active IL-6 signaling
326 in the liver. This result is consistent with the hepato-splanchnic removal of IL-6 previously
327 described in exercising human (12). IL-6 increases endogenous glucose production during
328 exercise by an unknown, hormone-independent mechanism (11) and because we found an
329 active signaling in the liver, we hypothesized that IL-6 could act directly on hepatic tissue to
330 modulate HGP during prolonged and intense exercise.

331 As expected in an exhaustive running exercise, plasma glucose was slightly
332 lower at the end of exercise and gluco-regulatory hormones measurement showed a drop in
333 plasma insulin/glucagon ratio and an increase in plasma corticosterone at the end of exercise.
334 There was a 86% decrease in plantaris muscle glycogen levels. We found both a profound
335 decrease in hepatic glycogen content (95%) and a substantial increase in liver key
336 gluconeogenic enzymes mRNA, PEPCK and G6Pase. PEPCK is regulated at the
337 transcriptional level and increased mRNA levels during exercise are associated to increased
338 enzyme activity (14, 44). Interestingly, we found a concomitant and substantial increase in
339 PGC-1 α mRNA, a key regulator of fasting-induced hepatic GNG (46). Taken together, these
340 data confirm that our model of long lasting exercise is a challenge for glucose homeostasis
341 and that HGP is very dependent on GNG at the end of exercise. Muscle IL-6 gene
342 transcription starts early during exercise, but IL-6 release in the plasma is delayed (17, 25).
343 Because circulating IL-6 concentration peaks at the end of exercise when GNG becomes
344 crucial to maintain plasma glucose levels, and because this pathway is regulated at the

345 transcriptional level, we hypothesized that IL-6 could favor HGP by modulating
346 gluconeogenic genes mRNA levels, particularly PEPCK.

347 PEPCK control during exercise is complex and in order to understand if IL-6 is able to
348 acutely modulate gluconeogenic genes independent of exercise-related factors, we injected 2
349 different doses of recombinant IL-6 to inactive rats and explored hepatic PEPCK and G6Pase
350 mRNA levels one hour after injection. Injected doses were calculated to mimic exercise-
351 induced plasma concentration. To avoid post-absorptive GNG blockade by insulin, animals
352 were fasted overnight. Injections resulted in a rise in circulating IL-6, with final plasma
353 concentrations flanking concentrations observed at the end of exhaustive running exercise. As
354 expected, we found a dose dependent increase in hepatic SOCS-3 and IL-6R mRNA levels
355 reflecting IL-6 signaling activation in the liver. This stimulation induced a dose dependent
356 increase in PEPCK mRNA levels, consistent with in vitro results (5). During food
357 deprivation, increased hepatic PEPCK mRNA levels result from both increased gene
358 transcription and mRNA stabilization through AMPc (16). In the present work, we did not
359 measure gene transcription, however the important and rapid increase in mRNA levels
360 suggests that PEPCK transcripts are stabilized in response exercise and IL-6 injection. No
361 changes in PGC-1 α and G6Pase mRNA abundance were found. Since G6Pase catalyses the
362 last step of glucose production, allowing glucose to leave the hepatocyte, this enzyme is
363 common to both GNG and glycogenolysis. In our experiment, rats were fasted overnight
364 (approximately 14 hours). In this situation, gluconeogenesis accounts for only 40-50% of
365 hepatic glucose output (24, 27), whereas glycogenolysis is fully induced after 4 to 6 hours
366 food deprivation (6). Therefore, a full activation of PEPCK gene is not expected, whereas
367 G6Pase was probably already activated, and no additive effect of IL-6 injection was found.
368 An increase in liver PGC-1 α mRNA levels at the end of 1 hour exercise as been previously
369 reported (19) and we show here that this is a transient rise. Whether newly transcribed PGC-

370 1α is involved in PEPCK transcription during exercise is not known and need to be explored,
371 however our results show that IL-6 effect on PEPCK gene is independent of any effect on
372 PGC-1 α mRNA levels. Plasma insulin/glucagon ratio and corticosterone were unchanged in
373 IL-6 injected rats, suggesting that gluco-regulatory hormones do not mediate IL-6 effect. We
374 found that PEPCK mRNA levels correlated with SOCS-3 and IL-6R mRNA levels in control
375 and exercised rats and this correlation was stronger in IL-6 injected rats, further suggesting a
376 co-regulation by IL-6 (Fig 8A and 8B). Differences observed between the two situations may
377 be due to the fact that hormonal factors are the main regulators of hepatic PEPCK mRNA
378 levels during exercise.

379 Independently of the debate concerning IL-6 and insulin sensitivity, the effect of IL-6
380 on HGP is still controversial. This cytokine has been described in vivo to whether activate
381 (40, 41) or inhibit HGP (7, 21). These differences could be explained by very different
382 environment in term of gluco-regulatory hormones, cytokines, gluconeogenic substrates, but
383 also by very different stimulation times (ranging from hours to days) and injected doses. Two
384 studies have investigated the effect of one ip injection of IL-6 in overnight fasted rodents, and
385 found no effect on PEPCK gene transcription (18, 34). The discrepancy between these results
386 and the increase in PEPCK mRNA we report could be explained by a substantially higher
387 hepatic IL-6 signaling activation in previous studies. Indeed, in the first study, authors
388 injected high IL-6 doses to mimic the same acute phase response as an acute inflammatory
389 agent (turpentine) and stimulation time was longer (18). In the second study (34), despite
390 similar fasting time, doses and stimulation time, the authors report a 30 fold increase in
391 hepatic SOCS-3 mRNA whereas we found a 12 or 4.2 fold increase in exercise and injection
392 studies respectively. Thus, it appears that acute effects of IL-6 on gluconeogenic genes are
393 very dependent on circulating concentrations. In our experiment, animals were fasted to
394 mimic exercise-induced variations in hormones and IL-6 was injected alone. Furthermore,

395 animals were sacrificed only 1h after injection to mimic the very transient IL-6 stimulation we
396 observed during exercise and plasma concentrations were similar to concentrations observed
397 in exhaustive exercise. Therefore, although we did not directly measure HGP or
398 gluconeogenic flux, our results are consistent with the positive effect of IL-6 on HGP during
399 prolonged exercise in human (11), and strongly suggest that IL-6 is able to directly modulate
400 liver PEPCK mRNA to increase HGP through GNG.

401 Plasma IL-6 concentrations obtained in our IL-6-injected animals reach the highest
402 levels described in human for intense and prolonged exercise (30). However, rats injected
403 with 3 or 10 $\mu\text{g}\cdot\text{kg}^{-1}$ recombinant IL-6 had only a 2 and 2.7 fold increase in hepatic PEPCK
404 mRNA (versus 5.5 at the end of prolonged intense exercise). This further confirms that
405 muscle-derived IL-6 cannot be considered a major regulator of glucose homeostasis during
406 exercise but rather a positive modulator of HGP, acutely supporting hormonal control in very
407 demanding exercises (11).

408

409 **Conclusion**

410 Prolonged intense exercise is a very challenging physiological situation for whole-
411 body glucose homeostasis. IL-6 has been shown to increase endogenous glucose production
412 during exercise in human through an unknown mechanism. Here we show that exhaustive
413 running exercise in rats induces an increase in plasma IL-6 and an active IL-6 signaling in the
414 liver. Because HGP was very dependent on GNG at the end of exercise, and GNG is regulated
415 at the transcriptional level, we tested IL-6 ability to modulate gluconeogenic genes mRNA
416 abundance in resting rats. We found a dose dependent increase in PEPCK mRNA levels in
417 response to IL-6 injection, with no changes in glucoregulatory hormones. Taken together, our
418 results suggest that muscle-derived IL-6 could contribute increasing HGP by directly up-
419 regulating PEPCK during intense and prolonged exercise.

420

421 Bibliography

- 422 1. **Akira S.** IL-6-regulated transcription factors. *Int J Biochem Cell Biol* 29: 1401-1418,
423 1997.
- 424 2. **Akira S, Isshiki H, Sugita T, Tanabe O, Kinoshita S, Nishio Y, Nakajima T,**
425 **Hirano T, and Kishimoto T.** A nuclear factor for IL-6 expression (NF-IL6) is a member of a
426 C/EBP family. *Embo J* 9: 1897-1906, 1990.
- 427 3. **Banzet S, Koulmann N, Sanchez H, Serrurier B, Peinnequin A, Alonso A, and**
428 **Bigard X.** Contraction-induced interleukin-6 transcription in rat slow-type muscle is partly
429 dependent on calcineurin activation. *J Cell Physiol* 210: 596-601, 2007.
- 430 4. **Bergmeyer H.** *Methods of enzymatic analysis.* New york: Academics, 1983.
- 431 5. **Blumberg D, Hochwald S, Brennan MF, and Burt M.** Interleukin-6 stimulates
432 gluconeogenesis in primary cultures of rat hepatocytes. *Metabolism* 44: 145-146, 1995.
- 433 6. **Cahill GF.** Fuel Metabolism in Starvation. *Annual Review of Nutrition* 26: 1-22, 2006.
- 434 7. **Christ B, Nath A, and Jungermann K.** Mechanism of the impairment of the
435 glucagon-stimulated phosphoenolpyruvate carboxykinase gene expression by interleukin-6 in
436 rat hepatocytes: inhibition of the increase in cyclic 3',5' adenosine monophosphate and the
437 downstream cyclic 3',5' adenosine monophosphate action. *Hepatology* 26: 73-80, 1997.
- 438 8. **Coker RH and Kjaer M.** Glucoregulation during exercise : the role of the
439 neuroendocrine system. *Sports Med* 35: 575-583, 2005.
- 440 9. **Coker RH, Simonsen L, Bulow J, Wasserman DH, and Kjaer M.** Stimulation of
441 splanchnic glucose production during exercise in humans contains a glucagon-independent
442 component. *Am J Physiol Endocrinol Metab* 280: E918-927, 2001.
- 443 10. **Dohm GL, Kasperek GJ, and Barakat HA.** Time course of changes in
444 gluconeogenic enzyme activities during exercise and recovery. *Am J Physiol* 249: E6-11,
445 1985.
- 446 11. **Febbraio MA, Hiscock N, Sacchetti M, Fischer CP, and Pedersen BK.** Interleukin-
447 6 is a novel factor mediating glucose homeostasis during skeletal muscle contraction.
448 *Diabetes* 53: 1643-1648, 2004.
- 449 12. **Febbraio MA, Ott P, Nielsen HB, Steensberg A, Keller C, Krstrup P, Secher**
450 **NH, and Pedersen BK.** Hepatosplanchnic clearance of interleukin-6 in humans during
451 exercise. *Am J Physiol Endocrinol Metab* 285: E397-402, 2003.
- 452 13. **Febbraio MA and Pedersen BK.** Muscle-derived interleukin-6: mechanisms for
453 activation and possible biological roles. *Faseb J* 16: 1335-1347, 2002.
- 454 14. **Friedman JE.** Role of glucocorticoids in activation of hepatic PEPCK gene
455 transcription during exercise. *Am J Physiol* 266: E560-566, 1994.
- 456 15. **Fujimoto M and Naka T.** Regulation of cytokine signaling by SOCS family
457 molecules. *Trends Immunol* 24: 659-666, 2003.
- 458 16. **Hanson RW and Reshef L.** Regulation of phosphoenolpyruvate carboxykinase
459 (GTP) gene expression. *Annu Rev Biochem* 66: 581-611, 1997.
- 460 17. **Helge JW, Stallknecht B, Pedersen BK, Galbo H, Kiens B, and Richter EA.** The
461 effect of graded exercise on IL-6 release and glucose uptake in human skeletal muscle. *J*
462 *Physiol* 546: 299-305, 2003.
- 463 18. **Hill M and McCallum R.** Altered transcriptional regulation of phosphoenolpyruvate
464 carboxykinase in rats following endotoxin treatment. *J Clin Invest* 88: 811-816, 1991.
- 465 19. **Hoene M, Lehmann R, Hennige AM, Pohl AK, Haring HU, Schleicher ED, and**
466 **Weigert C.** Acute regulation of metabolic genes and insulin receptor substrates in the liver of
467 mice by one single bout of treadmill exercise. *J Physiol* 587: 241-252, 2009.
- 468 20. **Howlett K, Febbraio M, and Hargreaves M.** Glucose production during strenuous
469 exercise in humans: role of epinephrine. *Am J Physiol* 276: E1130-1135, 1999.

- 470 21. **Inoue H, Ogawa W, Asakawa A, Okamoto Y, Nishizawa A, Matsumoto M,**
471 **Teshigawara K, Matsuki Y, Watanabe E, Hiramatsu R, Notohara K, Katayose K,**
472 **Okamura H, Kahn CR, Noda T, Takeda K, Akira S, Inui A, and Kasuga M.** Role of
473 hepatic STAT3 in brain-insulin action on hepatic glucose production. *Cell Metab* 3: 267-275,
474 2006.
- 475 22. **John-Alder HB, McAllister RM, and Terjung RL.** Reduced running endurance in
476 gluconeogenesis-inhibited rats. *Am J Physiol* 251: R137-142, 1986.
- 477 23. **Kamimura D, Ishihara K, and Hirano T.** IL-6 signal transduction and its
478 physiological roles: the signal orchestration model. *Rev Physiol Biochem Pharmacol* 149: 1-
479 38, 2003.
- 480 24. **Katz J and Tayek JA.** Gluconeogenesis and the Cori cycle in 12-, 20-, and 40-h-
481 fasted humans. *Am J Physiol Endocrinol Metab* 275: E537-542, 1998.
- 482 25. **Keller C, Steensberg A, Pilegaard H, Osada T, Saltin B, Pedersen BK, and**
483 **Neufer PD.** Transcriptional activation of the IL-6 gene in human contracting skeletal muscle:
484 influence of muscle glycogen content. *Faseb J* 15: 2748-2750, 2001.
- 485 26. **Kjaer M.** Hepatic metabolism during exercise. In: *Exercise metabolism*, edited by
486 Hargreaves MS, LL. Champaign, IL: human kinetics, 1998, p. 73-97.
- 487 27. **Landau BR, Wahren J, Chandramouli V, Schumann WC, Ekberg K, and Kalhan**
488 **SC.** Contributions of gluconeogenesis to glucose production in the fasted state. *J Clin Invest*
489 98: 378-385, 1996.
- 490 28. **Livak KJ and Schmittgen TD.** Analysis of relative gene expression data using real-
491 time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 25: 402-408, 2001.
- 492 29. **Nizielski SE, Arizmendi C, Shteyngarts AR, Farrell CJ, and Friedman JE.**
493 Involvement of transcription factor C/EBP-beta in stimulation of PEPCK gene expression
494 during exercise. *Am J Physiol* 270: R1005-1012, 1996.
- 495 30. **Ostrowski K, Rohde T, Zacho M, Asp S, and Pedersen BK.** Evidence that
496 interleukin-6 is produced in human skeletal muscle during prolonged running. *J Physiol* 508 (
497 Pt 3): 949-953, 1998.
- 498 31. **Park EA, Roesler WJ, Liu J, Klemm DJ, Gurney AL, Thatcher JD, Shuman J,**
499 **Friedman A, and Hanson RW.** The role of the CCAAT/enhancer-binding protein in the
500 transcriptional regulation of the gene for phosphoenolpyruvate carboxykinase (GTP). *Mol*
501 *Cell Biol* 10: 6264-6272, 1990.
- 502 32. **Pedersen BK and Febbraio MA.** Muscle as an endocrine organ: focus on muscle-
503 derived interleukin-6. *Physiol Rev* 88: 1379-1406, 2008.
- 504 33. **Puigserver P and Spiegelman BM.** Peroxisome proliferator-activated receptor-
505 gamma coactivator 1 alpha (PGC-1 alpha): transcriptional coactivator and metabolic
506 regulator. *Endocr Rev* 24: 78-90, 2003.
- 507 34. **Ramadoss P, Unger-Smith NE, Lam FS, and Hollenberg AN.** STAT3 targets the
508 regulatory regions of gluconeogenic genes in vivo. *Mol Endocrinol* 23: 827-837, 2009.
- 509 35. **Ramji DP and Foka P.** CCAAT/enhancer-binding proteins: structure, function and
510 regulation. *Biochem J* 365: 561-575, 2002.
- 511 36. **Saito M, Yoshida K, Hibi M, Taga T, and Kishimoto T.** Molecular cloning of a
512 murine IL-6 receptor-associated signal transducer, gp130, and its regulated expression in
513 vivo. *J Immunol* 148: 4066-4071, 1992.
- 514 37. **Sherman WM.** Metabolism of sugars and physical performance. *Am J Clin Nutr* 62:
515 228S-241S, 1995.
- 516 38. **Steensberg A, Febbraio MA, Osada T, Schjerling P, van Hall G, Saltin B, and**
517 **Pedersen BK.** Interleukin-6 production in contracting human skeletal muscle is influenced by
518 pre-exercise muscle glycogen content. *J Physiol* 537: 633-639, 2001.

- 519 39. **Steensberg A, van Hall G, Osada T, Sacchetti M, Saltin B, and Klarlund**
520 **Pedersen B.** Production of interleukin-6 in contracting human skeletal muscles can account
521 for the exercise-induced increase in plasma interleukin-6. *J Physiol* 529 Pt 1: 237-242, 2000.
- 522 40. **Stith RD and Luo J.** Endocrine and carbohydrate responses to interleukin-6 in vivo.
523 *Circ Shock* 44: 210-215, 1994.
- 524 41. **Tsigos C, Papanicolaou DA, Kyrou I, Defensor R, Mitsiadis CS, and Chrousos**
525 **GP.** Dose-dependent effects of recombinant human interleukin-6 on glucose regulation. *J*
526 *Clin Endocrinol Metab* 82: 4167-4170, 1997.
- 527 42. **Turcotte LP, Rovner AS, Roark RR, and Brooks GA.** Glucose kinetics in
528 gluconeogenesis-inhibited rats during rest and exercise. *Am J Physiol* 258: E203-211, 1990.
- 529 43. **Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, and**
530 **Speleman F.** Accurate normalization of real-time quantitative RT-PCR data by geometric
531 averaging of multiple internal control genes. *Genome Biol* 3: RESEARCH0034, 2002.
- 532 44. **Wasserman DH and Cherrington AD.** Hepatic fuel metabolism during muscular
533 work: role and regulation. *Am J Physiol* 260: E811-824, 1991.
- 534 45. **Wasserman DH, Spalding JA, Lacy DB, Colburn CA, Goldstein RE, and**
535 **Cherrington AD.** Glucagon is a primary controller of hepatic glycogenolysis and
536 gluconeogenesis during muscular work. *Am J Physiol* 257: E108-117, 1989.
- 537 46. **Yoon JC, Puigserver P, Chen G, Donovan J, Wu Z, Rhee J, Adelmant G, Stafford**
538 **J, Kahn CR, Granner DK, Newgard CB, and Spiegelman BM.** Control of hepatic
539 gluconeogenesis through the transcriptional coactivator PGC-1. *Nature* 413: 131-138, 2001.
- 540
541
542
543

544

545 **Legends**

546

547 **Figure 1: Liver carbohydrate metabolism in response to exhaustive exercise**

548 Liver glycogen content (A), PEPCK (B), G6P (C) and PGC-1 α (D) mRNA levels were
549 measured in resting rats (Rest), at the end of exercise (Ex), 2 hours (2h) and 6 hours (6h) after
550 the end of exercise.

551 * : significantly different from the other groups, $P < 0.001$.

552

553 **Figure 2: Glucoregulatory hormones in response to exhaustive exercise**

554 Plasma insulin and glucagons ratio (A) and corticosterone levels (B) were measured in resting
555 rats (Rest), at the end of exercise (Ex), 2 hours (2h) and 6 hours (6h) after the end of exercise.

556 # : significantly different from Rest group, $P < 0.05$; ## : significantly different from Rest
557 group, $P < 0.01$.

558

559 **Figure 3: Liver IL-6 signaling in response to exhaustive exercise**

560 Plasma IL-6 protein level (A), liver total and phospho STAT-3 (B) and C/EBP β (C) were
561 measured and phospho/total ratio was calculated in resting rats (Rest), at the end of exercise
562 (Ex), 2 hours (2h) and 6 hours (6h) after the end of exercise.

563 * : significantly different from the other groups, $P < 0.05$; ** : significantly different from the
564 other groups, $P < 0.01$.

565

566 **Figure 4: Liver IL-6-inductible genes mRNA levels in response to exhaustive exercise**

567 Liver SOCS-3 (A), C/EBP- β (B) and IL-6R (C) mRNA levels were measured in resting rats
568 (Rest), at the end of exercise (Ex), 2 hours (2h) and 6 hours (6h) after the end of exercise.

569 * : significantly different from the other groups, $P < 0.001$; # : significantly different from Rest
570 group, $P < 0.05$.

571

572 **Figure 5: Plasma rh IL-6 protein levels, liver SOCS-3 and IL-6R mRNA levels in 573 response to recombinant IL-6 injection.**

574 Plasma recombinant human IL-6 protein levels (A) Liver SOCS-3 (B) and IL-6R (C) mRNA
575 levels were measured in resting fasted rats 1 hour after recombinant IL-6 injection (3 or
576 10 $\mu\text{g.kg bodyweight}^{-1}$) or an equal volume of PBS buffer (Cont).

577 * : significantly different from Cont group, $P < 0.05$; # : significantly different from 3 $\mu\text{g.kg}^{-1}$
578 group, $P < 0.05$.

579

580 **Figure 6: Liver gluconeogenic genes mRNA levels in response to recombinant IL-6 581 injection.**





582 Liver PEPCK (A), G6P (B), PGC-1 α (C) mRNA levels were measured in resting fasted rats 1
583 hour after recombinant IL-6 injection (3 or 10 $\mu\text{g.kg bodyweight}^{-1}$) or an equal volume of PBS
584 buffer (Cont).



585 * : significantly different from Cont group, $P < 0.05$; ** : significantly different from Cont
586 group, $P < 0.01$.


587




588 **Figure 7: Association between IL-6 responsive genes and PEPCK mRNA in liver in 589 response to exercise and to recombinant IL-6 injection.**




590 Scatterplot showing SOCS-3 or IL-6R against PEPCK mRNA in liver from Cont and Ex
591 groups in exercise experiment (*A* and *B*), and in response to injection of 3 or 10 μ g.kg
592 bodyweight⁻¹ recombinant IL-6, or an equal volume of PBS buffer (*C* and *D*)























Gene	Accession number	Primers Sequences	Primers (μ M)	MgCl ₂ (mM)	Annealing temperature (°C)
Actine β	NM_031144 (816-910)	F : TCAGGTCATCACTATCGGCAATG R : TTTCATGGATGCCACAGGATTC	0,6	4	58
ARBP	NM_022402 (659-732)	F : CCTGCACACTCGCTTCCTAGAG R : CAACAGTCGGGTAGCCAATCTG	0,5	4	57
CycA	M19533 (381-507)	F : TATCTGCACTGCCAAGACTGAGTG R : CTTCTTGCTGGTCTTGCCATTCC	0,4	4	58
GAPDH	NM_017008 (1556-1633)	F : CCAATGTATCCGTTGTGGATCTGAC R : GCTTCACCACCTTCTTGATGTCATC	0,5	4	57
C/EBP- β	NM_024125 (656-760)	F : TGGACAAGCTGAGCGACGAG R : TGTGCTGCGTCTCCAGGTTG	0,4	4	59
PEPCK	NM_198780 (846-952)	F : TACGGTGGGAACACTGCTTG R : ATGCCCAGGATCAGCATGTG	0,4	4	59
G6PC	NM_013098 (173-304)	F : CCATCTGGTTCCACATTCAAGAG R : CAGTGTCAGGACCCACCAATAC	0,4	4	59
HPRT	S79292 (211-333)	F : CTCATGGACTGATTATGGCAGGAC R : GCAGGTCAGCAAAGAACTTATAGCC	0,4	4	60
IL-6 R	NM_017020 (696-812)	F : AAGCAGGTCCAGCCACAATGTAG R : CCAACTGACTTTGAGCCAACGAG	0,4	5	60
IL-6	E02522 (532-610)	F : TCCTACCCCAACTCCAATGCTC R : TTGGATGGTCTTGGTCCTTAGCC	0,4	3	65
SOCS-3	AF075383 (581-679)	F : CCTCCAGCATCTTTGTGCGGAAGAC R : TACTGGTCCAGGAACTCCCGAATG	0,4	4	65
PGC-1 α	NM_031347 (2249-2364)	F : ACGCAGGTCGAATGAACTGAC R : TGGTGAAGCAGGGTCAAAAATC	0,4	4	53

Table 1